

# ISMP's List of *Error-Prone Abbreviations, Symbols, and Dose Designations*

The abbreviations, symbols, and dose designations found in this table have been reported to ISMP through the ISMP National Medication Errors Reporting Program (ISMP MERP) as being frequently misinterpreted and involved in harmful medication errors. They should **NEVER** be used when commu-

nicating medical information. This includes internal communications, telephone/verbal prescriptions, computer-generated labels, labels for drug storage bins, medication administration records, as well as pharmacy and prescriber computer order entry screens.

Abbreviations	Intended Meaning	Misinterpretation	Correction
µg	Microgram	Mistaken as "mg"	Use "mcg"
AD, AS, AU	Right ear, left ear, each ear	Mistaken as OD, OS, OU (right eye, left eye, each eye)	Use "right ear," "left ear," or "each ear"
OD, OS, OU	Right eye, left eye, each eye	Mistaken as AD, AS, AU (right ear, left ear, each ear)	Use "right eye," "left eye," or "each eye"
BT	Bedtime	Mistaken as "BID" (twice daily)	Use "bedtime"
cc	Cubic centimeters	Mistaken as "u" (units)	Use "mL"
D/C	Discharge or discontinue	Premature discontinuation of medications if D/C (intended to mean "discharge") has been misinterpreted as "discontinued" when followed by a list of discharge medications	Use "discharge" and "discontinue"
IJ	Injection	Mistaken as "IV" or "intrajugular"	Use "injection"
IN	Intranasal	Mistaken as "IM" or "IV"	Use "intranasal" or "NAS"
HS	Half-strength	Mistaken as bedtime	Use "half-strength" or "bedtime"
hs	At bedtime, hours of sleep	Mistaken as half-strength	
IU**	International unit	Mistaken as IV (intravenous) or 10 (ten)	Use "units"
o.d. or OD	Once daily	Mistaken as "right eye" (OD-oculus dexter), leading to oral liquid medications administered in the eye	Use "daily"
OJ	Orange juice	Mistaken as OD or OS (right or left eye); drugs meant to be diluted in orange juice may be given in the eye	Use "orange juice"
Per os	By mouth, orally	The "os" can be mistaken as "left eye" (OS-oculus sinister)	Use "PO," "by mouth," or "orally"
q.d. or QD**	Every day	Mistaken as q.i.d., especially if the period after the "q" or the tail of the "q" is misunderstood as an "i"	Use "daily"
qhs	Nightly at bedtime	Mistaken as "qhr" or every hour	Use "nightly"
qn	Nightly or at bedtime	Mistaken as "qh" (every hour)	Use "nightly" or "at bedtime"
q.o.d. or QOD**	Every other day	Mistaken as "q.d." (daily) or "q.i.d. (four times daily) if the "o" is poorly written	Use "every other day"
q1d	Daily	Mistaken as q.i.d. (four times daily)	Use "daily"
q6PM, etc.	Every evening at 6 PM	Mistaken as every 6 hours	Use "daily at 6 PM" or "6 PM daily"
SC, SQ, sub q	Subcutaneous	SC mistaken as SL (sublingual); SQ mistaken as "5 every;" the "q" in "sub q" has been mistaken as "every" (e.g., a heparin dose ordered "sub q 2 hours before surgery" misunderstood as every 2 hours before surgery)	Use "subcut" or "subcutaneously"
ss	Sliding scale (insulin) or ½ (apothecary)	Mistaken as "55"	Spell out "sliding scale;" use "one-half" or "½"
SSRI	Sliding scale regular insulin	Mistaken as selective-serotonin reuptake inhibitor	Spell out "sliding scale (insulin)"
SSI	Sliding scale insulin	Mistaken as Strong Solution of Iodine (Lugol's)	
i/d	One daily	Mistaken as "tid"	Use "1 daily"
TIW or tiw	3 times a week	Mistaken as "3 times a day" or "twice in a week"	Use "3 times weekly"
U or u**	Unit	Mistaken as the number 0 or 4, causing a 10-fold overdose or greater (e.g., 4U seen as "40" or 4u seen as "44"); mistaken as "cc" so dose given in volume instead of units (e.g., 4u seen as 4cc)	Use "unit"
UD	As directed ("ut dictum")	Mistaken as unit dose (e.g., diltiazem 125 mg IV infusion "UD" misinterpreted as meaning to give the entire infusion as a unit [bolus] dose)	Use "as directed"
Dose Designations and Other Information	Intended Meaning	Misinterpretation	Correction
Trailing zero after decimal point (e.g., 1.0 mg)**	1 mg	Mistaken as 10 mg if the decimal point is not seen	Do not use trailing zeros for doses expressed in whole numbers
"Naked" decimal point (e.g., .5 mg)**	0.5 mg	Mistaken as 5 mg if the decimal point is not seen	Use zero before a decimal point when the dose is less than a whole unit
Abbreviations such as mg. or mL. with a period following the abbreviation	mg mL	The period is unnecessary and could be mistaken as the number 1 if written poorly	Use mg, mL, etc. without a terminal period

**ISMP's List of Error-Prone Abbreviations, Symbols, and Dose Designations** (continued)

Dose Designations and Other Information	Intended Meaning	Misinterpretation	Correction
Drug name and dose run together (especially problematic for drug names that end in "I" such as Inderal40 mg; Tegretol300 mg)	Inderal 40 mg Tegretol 300 mg	Mistaken as Inderal 140 mg Mistaken as Tegretol 1300 mg	Place adequate space between the drug name, dose, and unit of measure
Numerical dose and unit of measure run together (e.g., 10mg, 100mL)	10 mg 100 mL	The "m" is sometimes mistaken as a zero or two zeros, risking a 10- to 100-fold overdose	Place adequate space between the dose and unit of measure
Large doses without properly placed commas (e.g., 100000 units; 1000000 units)	100,000 units 1,000,000 units	100000 has been mistaken as 10,000 or 1,000,000; 1000000 has been mistaken as 100,000	Use commas for dosing units at or above 1,000, or use words such as 100 "thousand" or 1 "million" to improve readability
Drug Name Abbreviations	Intended Meaning	Misinterpretation	Correction
To avoid confusion, do not abbreviate drug names when communicating medical information. Examples of drug name abbreviations involved in medication errors include:			
<b>APAP</b>	acetaminophen	Not recognized as acetaminophen	Use complete drug name
<b>ARA A</b>	vidarabine	Mistaken as cytarabine (ARA C)	Use complete drug name
<b>AZT</b>	zidovudine (Retrovir)	Mistaken as azathioprine or aztreonam	Use complete drug name
<b>CPZ</b>	Compazine (prochlorperazine)	Mistaken as chlorpromazine	Use complete drug name
<b>DPT</b>	Demerol-Phenergan-Thorazine	Mistaken as diphtheria-pertussis-tetanus (vaccine)	Use complete drug name
<b>DTO</b>	Diluted tincture of opium, or deodorized tincture of opium (Paregoric)	Mistaken as tincture of opium	Use complete drug name
<b>HCl</b>	hydrochloric acid or hydrochloride	Mistaken as potassium chloride (The "H" is misinterpreted as "K")	Use complete drug name unless expressed as a salt of a drug
<b>HCT</b>	hydrocortisone	Mistaken as hydrochlorothiazide	Use complete drug name
<b>HCTZ</b>	hydrochlorothiazide	Mistaken as hydrocortisone (seen as HCT250 mg)	Use complete drug name
<b>MgSO4**</b>	magnesium sulfate	Mistaken as morphine sulfate	Use complete drug name
<b>MS, MSO4**</b>	morphine sulfate	Mistaken as magnesium sulfate	Use complete drug name
<b>MTX</b>	methotrexate	Mistaken as mitoxantrone	Use complete drug name
<b>PCA</b>	procainamide	Mistaken as patient controlled analgesia	Use complete drug name
<b>PTU</b>	propylthiouracil	Mistaken as mercaptopurine	Use complete drug name
<b>T3</b>	Tylenol with codeine No. 3	Mistaken as liothyronine	Use complete drug name
<b>TAC</b>	triamcinolone	Mistaken as tetracaine, Adrenalin, cocaine	Use complete drug name
<b>TNK</b>	TNKase	Mistaken as "TPA"	Use complete drug name
<b>ZnSO4</b>	zinc sulfate	Mistaken as morphine sulfate	Use complete drug name
Stemmed Drug Names	Intended Meaning	Misinterpretation	Correction
<b>"Nitro" drip</b>	nitroglycerin infusion	Mistaken as sodium nitroprusside infusion	Use complete drug name
<b>"NorfloX"</b>	norfloxacin	Mistaken as Norflex	Use complete drug name
<b>"IV Vanc"</b>	intravenous vancomycin	Mistaken as Invanz	Use complete drug name
Symbols	Intended Meaning	Misinterpretation	Correction
$\text{̄}$	Dram	Symbol for dram mistaken as "3"	Use the metric system
$\text{m̄}$	Minim	Symbol for minim mistaken as "mL"	
<b>x3d</b>	For three days	Mistaken as "3 doses"	Use "for three days"
<b>&gt; and &lt;</b>	Greater than and less than	Mistaken as opposite of intended; mistakenly use incorrect symbol; "< 10" mistaken as "40"	Use "greater than" or "less than"
<b>/ (slash mark)</b>	Separates two doses or indicates "per"	Mistaken as the number 1 (e.g., "25 units/10 units" misread as "25 units and 10 units")	Use "per" rather than a slash mark to separate doses
<b>@</b>	At	Mistaken as "2"	Use "at"
<b>&amp;</b>	And	Mistaken as "2"	Use "and"
<b>+</b>	Plus or and	Mistaken as "4"	Use "and"
<b>o</b>	Hour	Mistaken as a zero (e.g., q2 <sup>o</sup> seen as q 20)	Use "hr," "h," or "hour"
<b>Φ or ∅</b>	zero, null sign	Mistaken as numerals 4, 6, 8, and 9	Use 0 or zero, or describe intent using whole words

\*\*These abbreviations are included on The Joint Commission's "minimum list" of dangerous abbreviations, acronyms, and symbols that must be included on an organization's "Do Not Use" list, effective January 1, 2004. Visit [www.jointcommission.org](http://www.jointcommission.org) for more information about this Joint Commission requirement.